

Mesoscale Eddy Formation and Shock Features Associated with a Coastally Trapped Disturbance

Stephen D. Burk and William T. Thompson

Naval Research Laboratory

Monterey, California

burk@nrlmry.navy.mil

Abstract- Synoptic maps for 28-29 August 2002 indicate typical near-surface northerly flow along coastal California, while Goes-10 animated satellite imagery from this period show a stratus-capped, coastally trapped disturbance (CTD) propagating northward against the prevailing flow. A pronounced hydraulic-jump-like ('shock') feature develops in the cloud field south of Cape Mendocino (CM) and angles linearly away from the coast. Over the next ~6 hours both the CTD and the shock feature move northward and, as they approach the Cape, the leading edge of the CTD clouds roll-up into a visually striking cyclonic mesoscale eddy—with the shock feature being wrapped into the eddy. The CTD rather abruptly stalls and fails to round CM. Further, a second cyclonic mesoscale eddy is observed to form SW of Point Arena.

Using a mesoscale model, many of the observed features of this case are forecast, including: (a) the cloud-filled, northward propagating CTD; (b) the development, linear structure, and orientation of the oblique shock feature; (c) the CTD rounds Point Arena, but fails to round CM; and (d) the formation of cyclonic mesoscale eddies near CM and Point Arena. The model fidelity permits us to analyze the dynamics associated with the shock feature and eddy formation.

I. INTRODUCTION

On 28 August 2002, an interesting sequence of events took place in association with a coastally trapped disturbance (CTD) as it propagated along the coast of northern California. The CTD (as evidenced by a sharply defined, narrow tongue of marine stratus cloud) propagated northward against the prevailing northwesterly flow, arrived in Monterey Bay at 11:30 UTC, San Francisco Bay at 14:30 UTC, and reached Pt. Reyes at 16:30 UTC 28 August 2002. As the CTD rounded Pt. Arena at about 18:30 UTC, a dramatic, well-defined hydraulic-jump-like feature (hereafter termed a 'shock') developed north of the CTD and angled away from the coast to the southwest. Because the summertime marine boundary layer (MBL) in

this coastal region often is close to saturation, an abrupt increase in MBL depth associated with a shock feature sometimes can generate recognizable, and in some cases very eye-catching, features within the cloud field [1]. Over the next 3-4 h, the CTD continued to propagate to the north and then rolled into a striking mesoscale eddy leeward of Cape Mendocino (CM) and another eddy formed off of Pt. Arena.

Although a considerable body of literature recently has been devoted to the propagation phase of CTD events, very little study has been directed towards the CTD decay phase wherein mesoscale eddy formation is not unusual. It has previously been noted [2] that mesoscale eddies can be generated in association with CTD's and [2] presented satellite imagery from May 1982 of a CTD along coastal California bearing similarity to the August 2002 case we analyze here. There are instances, however, when CTD's propagate the full length of the U. S. west coast, opposite to prevailing northerly flow, without developing mesoscale eddies.

In [3] a shallow-water model was used to investigate the interaction of an artificially created CTD with transcritical flow in the vicinity of coastal topographic bends. Shock features develop at the southern end of an expansion fan in that study and, as the CTD propagates north along the coast, a strong lateral shear layer develops. Once CTD propagation is halted, the shear layer destabilizes and rapidly rolls up to form a vortex. Many of these same features occur in our mesoscale forecast presented here. However, the origin of the supercritical flow leeward of CM in our study is not solely due to flow acceleration in an expansion fan around CM. Supercritical flow also is produced by downslope acceleration in the lee of CM. Nonhydrostatic mesoscale model simulations of this period using the Coupled Ocean-Atmosphere Mesoscale Prediction System

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 01 SEP 2003		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Mesoscale Eddy Formation and Shock Features Associated with a Coastally Trapped Disturbance				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Monterey, California 93943				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002146. Oceans 2003 MTS/IEEE Conference, held in San Diego, California on September 22-26, 2003. U.S. Government or Federal Purpose Rights License., The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 8	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

(COAMPSTM)¹ reproduce features of the event seen in satellite images and surface observations.

II. MODELING ASPECTS

The COAMPS mesoscale model used in the present study is described in [4]. The model is nonhydrostatic and uses multiple nests having different horizontal resolution. It features a full suite of physical parameterizations, including a level 2.5 turbulence parameterization, radiation, and cloud microphysics schemes. Surface fluxes and surface stress are computed from the Louis scheme. Data assimilation is accomplished using a multivariate optimal interpolation (MVOI) approach.




Fig. 1 COAMPS nested grid structure used in this study.

The COAMPS model simulation of this event utilizes the grid structure shown in Fig 1. As indicated, the horizontal resolution of the nests is 27 km, 9 km, and 3 km. Three 12 h data assimilation forecasts were performed prior to the 12 h forecast beginning at 1200 UTC 28 August containing the period of interest.

III. CASE DESCRIPTION

A. Synoptic Discussion

On 27 August 2002, a weak 500-hPa trough was located over the western U. S. with the axis extending from Montana to Central California. At sea level, a ridge over Oregon and Washington resulted in NE offshore flow over Northern California while an inverted thermal trough was located over the southern San Joaquin Valley and the southern deserts of

California. On 28 August, the trough at 500 hPa deepened significantly; the axis extended from Montana to ~100 km west of San Diego. The inverted thermal trough over California deepened as well and extended from just east of CM to South-Central Arizona. The ridge over the Pacific Northwest resulted in continued offshore flow over Northern California (the high in San Francisco was 29.4C; 11 C above the monthly mean for August 2002). On 29 August, the 500-hPa trough weakened substantially, as did the thermal trough at the surface over California.

B. Satellite Imagery

The mesoscale evolution of the event can be followed by a series of GOES10 satellite images from the afternoon of 28 August. Hereinafter, time will be specified in local time (LT=PDT=UTC-7). The image from 1300 LT 28 August 2002 (Fig. 2a) shows a cloud-free zone along the coast of Southern Oregon and Northern California. There is a large area of low clouds 50-75 km offshore extending along the coast over the domain of the image. The narrow cloud "tongue" that has rounded Pt. Arena is associated with the CTD.

West of the CTD at the edge of the cloud band is a series of wave clouds. These wave clouds are indicative of trapped gravity waves associated with a shock feature and are similar in appearance to wave clouds discussed in [1]. The wave clouds are distinct, having an albedo significantly larger than the surrounding low clouds.




Fig. 2 GOES10 images at (a) 2000 UTC [1300 LT] and, (b) 2200 UTC [1500 LT] 28 Aug 2002.

¹ COAMPS is a trademark of the Naval Research Laboratory.




Fig. 2 (b)




Fig. 3a-b

Fig. 3(a) 10-m streamlines and integrated liquid water from 12 h COAMPS forecast valid 1700 LT 28 Aug 2002 and, (b) 10-m vectors and shaded wind speed (m s^{-1}) at same time.

At 1500 LT (Fig 2b), a rather striking image shows that a pair of cyclonic eddies have formed due to the interaction of the prevailing NW coast-parallel flow and the southerly flow

associated with the CTD. The northern eddy is just in the lee of CM and is delineated by fine cloud filaments wrapping around the center while the southern eddy lies just to the SW of Pt. Arena. The southern eddy is less distinct than the northern eddy in individual satellite images, but can be seen clearly in cloud motions visualized by a satellite loop. These eddies are both $\sim 50\text{--}75$ km in diameter. The linear feature associated with the shock is somewhat less visible at 1500 LT, having been entangled in the northern eddy.

The switch to southerly flow at a given location along the coast precedes the arrival of the cloud tongue by several hours, as evidenced by comparing buoy and satellite observations. This same lag between wind shift and cloud arrival is discussed in [5] and modeled in [6] in conjunction with a June 1994 CTD. The air in this forerunning, wind-shift region is not purely associated with southerly flow and, hence, saturation is not immediately attained upon the switch to southerly flow.

C. Model Forecasts

Shown in Fig 3a are forecasts of 10 m elevation streamlines and integrated cloud liquid water (kg m^{-2} ; shaded) over a zoomed area of the inner nest. Figure 3a shows the model 12 h forecast valid 1700 LT 28 Aug. At this time the modeled zone of sharp wind shift associated with the shock has strengthened substantially and propagated to the north in agreement with the satellite images. Both of the model eddies are well defined and southerly flow extends to just south of CM in a narrow zone along the immediate coast in agreement with the satellite images. The cloud tongue associated with the CTD has rounded Pt Arena and, as in the satellite imagery, a clear "slot" is visible over the center of the southern eddy. The cloudy area to the west has advanced toward the coast. Note that the modeled integrated cloud water is much larger within the southerly flow associated with the CTD than in the cloud shield to the west, which is consistent with the higher albedo of these clouds in the satellite images.

Figure 3b shows model forecast wind vectors and isotachs (m s^{-1} ; gray shaded) at 10 m at the same time as Fig 3a. There is strong flow acceleration in the lee of CM, with abrupt deceleration along a line (shock) angling away from the coast to the SW. Another shock feature has formed in the blocked flow north of CM.

D. Comparison with surface observations

Time series of wind speed and direction and temperature have been constructed at locations corresponding to the positions of several data buoys along the coast. The buoys are located at Pt. Piedras Blancas, Monterey Bay, San Francisco Bay, Bodega Bay, Pt. Arena, and Eel River. Comparison of the wind direction observations with model results allows tracking of the observed and modeled progression of the




Fig 3 (b)

CTD up the coast. The comparison shows good agreement between the model and observations from Pt. Piedras Blancas, where the model and observations show a distinct shift to southerly flow at 0200 LT, to Bodega Bay, where the shift to southerly flow commences at 0700 LT in the observations and the model results. In the observations, propagation of the CTD slows north of Bodega Bay; however, it slows far too much at the surface in the model results. A phase lag in surface wind shift between the model and observations of ~4 h develops at Pt. Arena. However, the model does have tongue of southerly flow just above the surface layer that extends well ahead of the surface southerly flow. In particular, at 140 m the modeled southerly flow reaches Pt. Arena at 1100 LT,

only one hour later than in the surface observations. Hence, it appears that the model may be too slow in mixing this southerly flow to the surface.

IV. Shock Feature Interaction with the CTD and Eddy Roll-up

Compression jumps, which tend to form when coastal points or capes act to block and deflect approaching supercritical flow, often take on the linear or the gently curved appearance of a bow wave. Modeling the formation of atmospheric shocks when supercritical flow impacts a convex bend in the coastline has generally been quite idealized, often making use of the shallow water equations.

In a single layered fluid, supercritical flow occurs when the Froude number (Fr), a dimensionless quantity given by the ratio of the flow speed to the gravity wave phase speed, is greater than unity. The properties of free and forced trapped waves in an atmosphere that includes a stable layer above the capping inversion were investigated in [7] and the gravest-mode phase speed was found to be increased, and higher vertical modes supported, by the presence of the extended stable layer. In fact, for an atmospheric structure as displayed in [7; his Fig. 1], $c = [(\gamma_1 + \gamma_2)h]^{1/2}$ is found to be a relatively good approximation to the phase speed of the first trapped mode. Here $\gamma_1 = g\Delta\Theta/\langle\Theta\rangle$, $\gamma_2 = g(\delta\Theta/\Theta)$ with $\Delta\Theta$ being the sharp discontinuity at the MBL top, and $\delta\Theta$ being the potential temperature change across the extended stable layer above the capping inversion, and h the MBL depth. The phase speed and layer depth we use in our COAMPS Froude number computations is similar to that described above.

Because of this imprecision in specifying a meaningful Fr value (for stratified flow) as an indicator of regimes of flow transition, we examine regions of sharp horizontal gradients in Fr within which the $Fr = 1$ line is embedded, rather than simply focusing on the $Fr=1$ line itself. This is illustrated in Fig. 4, which displays a 12h forecast of model computed Fr valid at 1700 LT 28 Aug 2002. Maximum Fr values greater than 1.6 occur leeward (south) of CM. This high Fr region is created by two processes; namely, flow acceleration down the lee slope of CM associated with a mountain gravity wave and a low-level cornering flow (expansion fan). Also, a local maximum in Fr occurs adjacent to

the coast south of Point Arena that is not due to an expansion fan but rather the strong,

(a)




Fig. 4 COAMPS forecast Froude number at same time as Fig. 3

southerly flow associated with the propagating CTD that has entered this region.

Most noteworthy in Fig. 4 is the region of strong horizontal Fr-gradient, which encompasses the $Fr = 1$ contour that angles away from the coast at the southern end of the CM expansion fan. There also is a pronounced Fr gradient north of CM that is indicative of a shock feature and, indeed, there is evidence of this feature in the satellite cloud imagery as well, albeit somewhat more subtle.

Figure 5 depicts the COAMPS forecast of the 0.05 g kg^{-1} isosurface of cloud liquid water, along with 10-m wind speed (m s^{-1}) and streamlines at 1700 LT. The linear cloud lines evident in the satellite imagery are oriented at $\sim 125^\circ$ from north (Fig. 2a), which is approximately the angle at which the modeled linear shock feature forms (Fig. 5). Also, the tight gradient associated with the modeled shock extends over a length of $\sim 100 \text{ km}$, which is approximately that of the observed cloud lines.




Fig. 5 COAMPS 12h forecast 10-m streamlines, wind speed shaded (m s^{-1}), and cloud water isosurface (0.05 g kg^{-1}) at same time as Fig. 3

Although Fig. 5 shows that the model's wind field has captured the shock feature, the cloud features along the shock are poorly represented. Resolution greater than the 3 km inner nest is required to adequately describe the undular wave clouds associated with the shock. In agreement with the satellite image (Fig. 2a), however, the modeled cloud field (Fig. 5) does show a clear gap between the cloud line along the shock and the coastline. This gap in the clouds arises because the MBL shallows progressively as the expansion fan/ downslope flow approaches the coast and consequently the inversion capping the MBL drops below the lifting condensation level.

The modeled southerly flow in the MBL associated with the CTD is strong (10-m wind speed maximum $\sim 8 \text{ m s}^{-1}$) as it rounds Point Arena, however the CTD fails to round CM and instead, in agreement with the observations, wraps into two eddies. The dynamic characteristics of hydraulically transcritical flows [that is, flows having localized supercritical ($Fr > 1$) regions embedded within a domain that generally is subcritical ($Fr < 1$)] adjacent to a spatially varying coastline was investigated in [1]. While CTD's are able to propagate northward in the subcritical region of the base flow, they can be severely attenuated, or halted altogether, upon encountering a supercritical region. In our real data forecast, prior to the CTD's approach to Point Arena the model

computed Fr field has only a very small supercritical area in the lee of this Point, but a much stronger supercritical region (with $Fr > 2.0$) leeward of CM, with a sizeable region encompassed by the $Fr = 1.5$ contour. Consistent with [3], the CTD modeled here is able to propagate northward beyond Point Arena, but is halted at CM. The area encompassed by the $Fr = 1.5$ contour has shrunk considerably by 1700 LT 28 Aug. (Fig. 4), and is absent by 2100 LT on the 28th (not shown), although the flow in a small region in the immediate lee of the Cape remains supercritical.

The COAMPS divergence and absolute vorticity fields in the MBL (140 m) are shown in Fig. 6a-b at 1700 LT 28 August 2002. Strong convergence zones are evident (Fig. 6a) associated with the modeled shock features either side of CM, while there is divergence in the accelerating flow leeward of the Cape. Prior to 1300 LT (not shown) the convergence zone angling away from the coast is weak and unorganized.

Vorticity is clearly enhanced in association with the shock feature (Fig. 6b). Idealized flow over a uniform ridge was modeled in [8] and they investigated the mechanisms of vorticity production associated with a hydraulic-jump feature that develops leeward of the ridge. It is found in [8] that the vertical stretching term [i.e., $(\zeta+f)\nabla \cdot \mathbf{V}$] in the jump significantly amplifies the upstream relative vorticity. Based on the divergence field shown in Fig. 6a, which indicates strong convergence along the shock line, vorticity stretching also appears likely to be a significant factor in generating the vorticity enhancement seen along the shock in this study. Of course, potential vorticity (PV) is a conservative quantity; stretching and tilting of relative or absolute vorticity does not generate PV. However, there is generation of PV along the shock feature in this study associated with dissipative processes within the jump. This PV enhancement can best be seen in a vertical cross section across the jump.




Fig. 6 (a) COAMPS 12h forecast of divergence (10^{-5} s^{-1}) at 140 m valid 1700 LT 28 Aug 2002 and, (b) absolute vorticity (10^{-5} s^{-1}) at same height and time as (a).

Figure 7a shows the potential temperature (K) in the plane of cross section E-F (Fig. 6b), and Fig. 7b displays the PV in this same plane. Both Fig. 7 a-b are from a 12h COAMPS forecast valid 1700 LT 28 August 2002.

V. Discussion and Conclusions

In the present study, we investigate an event in which a CTD propagating northward along the coast past Point Arena encounters supercritical flow in the lee of CM. Satellite imagery shows that a compression shock forms south of CM, angling away from the coast. The shock feature propagates to the




Fig. 6 (b)

north for several hours and then becomes rolled up into a cyclonic eddy. The Naval Research Laboratory's nonhydrostatic COAMPS model is used to simulate this period. The model results compare quite favorably with satellite imagery and coastal buoy observations, provided allowance is made for the modeled CTD phase speed being too slow near Pt. Arena.

Numerous recent investigations (including field studies) have been conducted addressing the propagation phase of CTD's, but there has been relatively little study of either the initiation or dissipation phase of CTD's. The coastal processes present in this case study are not particularly rare; indeed transcritical conditions are found in the lee of points and capes in a mean summer (June-July) model average [9] and CTD's occur several times each year, often spawning cyclonic eddies in conjunction with cessation of propagation. The event described herein, however, features all of these processes and incorporates the highly nonlinear interaction among them, including the northward propagation of the shock feature, the sharp variations in boundary layer depth associated with both the shock feature and the leading edge of the CTD, and the strong horizontal shear associated with the NW flow offshore and the southerly flow in the CTD aiding development of the cyclonic eddies




Fig. 7 (a) COAMPS 12h forecast of potential temperature (K) in section E-F of fig. 6b valid 1700 LT 28 Aug 2002 and, (b) potential vorticity (PVU) in the same plane at the same time as (a).




Fig. 7 (b)

Both the observed and modeled CTD round Pt. Arena with apparent ease and show rather strong southerly flow north of this Point. However, both the observed and modeled CTD stall and roll-up into a mesoscale eddy upon approaching CM. Prior to this eddy roll up, vorticity and PV are both clearly enhanced across the shock feature. The strong low-level convergence along the leading edge of the shock and the jump in boundary layer depth are favorable for vorticity enhancement by stretching [8]. But, alone this process only produces a modest increase in vorticity along the length of the shock (Fig. 6b). However, as the CTD approaches the shock feature, the combined interaction of these jump features acts to strongly enhance the vorticity at the leading edge of the CTD (*note strong maximum south of CM along the coast in Fig. 6b*). And, this is the point when the leading edge of the CTD first begins to roll up into a mesoscale vortex.

The eddy that forms off of Pt. Arena does not appear to form in the same manner as just described for the northern eddy. The southern eddy forms along the strong shear line between the southerly flow rounding Pt. Arena and the background northerly flow. This eddy appears to grow by shear (barotropic) instability wherein the perturbation directly extracts energy from the mean shear. This is further indicated by the orientation of this eddy relative to the shear vector [see 10 for further discussion of this instability mechanism].

ACKNOWLEDGEMENTS

We gratefully acknowledge Prof. R. T. Williams for helpful discussions and Dr. Peter Guest for bringing this CTD event to our attention. This research was supported by the Office of Naval Research Program Element 0601153N.

References

- [1] Burk, S.D. and T. Haack, 2001: The dynamics of wave clouds upwind of coastal orography. *MWR.*, **128**, 1438-1455.
- [2] Dorman, C.E., 1985 : Evidence of Kelvin waves in California's marine layer and related energy generation. *MWR*, **113**, 827-839.
- [3] Rogerson, A. M., 1999: Transcritical flows in the coastal marine atmospheric boundary

- layer. *J. Atmos. Sci.*, **56**, 2761-2779.
- [4] Hodur, R. M., J. Pullen, J. Cummings, X. Hong, J.D. Doyle, P. Martin, and M.A. Rennick, 2002: The coupled ocean/atmospheric mesoscale prediction system (COAMPS). *Oceanography*, **15**, 88-89.
- [5] Ralph, F. M., L. Armi, J. M. Bane, C. Dorman, W. D. Neff, P. J. Neiman, W. Nuss, and P.O.G. Persson , 1998 : Observations and analysis of the 10-11 June coastally trapped disturbance. *MWR*, **126**, 2435-2465.
- [6] Thompson, W. T., T. Haack, J. D. Doyle, and S. D. Burk, 1997: A nonhydrostatic mesoscale simulation of the 10-11 June 1994 coastally trapped wind reversal. *MWR*, **125**, 3211-3230.
- [7] Samelson, R. M., 1999: The vertical structure of linear coastal-trapped disturbances. *MWR*, **127**, 201-213.
- [8] Epifanio, C.C., and D.R. Durran, 2002: Lee-vortex formation in free-slip stratified flow over ridges. II: Mechanisms of vorticity and PV production in nonlinear viscous wakes. *J. Atmos. Sci.*, **59**, 1166-1181.
- [9] -----, T. Holt, D.P. Rogers, and K. Edwards, 2000: Large-scale structure of the June-July 1996 marine boundary layer along California and Oregon. *MWR*, **128**, 1632-1652.
- [10] Pedlosky, J., 1979: *Geophysical Fluid Dynamics*. Springer-Verlag, 624 pp.

Corresponding author : Stephen D. Burk, Naval Research Laboratory, Marine Meteorology Division, 7 Grace Hopper Ave., Monterey, CA 93943-5502 : burk@nrlmry.navy.mil